



# CARE GUIDE

---

Tools and resources to support  
your treatment journey

# Introducing The TECENTRIQ Care Guide

We're committed to supporting you throughout your treatment journey. So we created this guide with your needs in mind to help you stay on track.

Inside this folder, you'll find these resources:



## **Understanding Your Disease**

You may already be familiar with the basics of your condition. In case you're just getting started, this resource can give you more background



## **Where to Find More Information and Support**

Finding the right source of support for your disease can be challenging. Here you can discover a list of organizations that you may find helpful



## **What to Know About TECENTRIQ + Avastin® (bevacizumab)**

This resource explains the clinical study results of TECENTRIQ + Avastin. You'll also find information on how TECENTRIQ works and how it's given



## **Options That Can Help You Afford Treatment**

Find out how Genentech may help you get your medicines, regardless of your insurance status



## **Learn About Potential Side Effects**

TECENTRIQ can cause serious side effects. Here you can find a list of common side effects of TECENTRIQ



## **Ways to Help Manage Side Effects**

Discover tips that can help you manage some of the common side effects of TECENTRIQ




## **My Medication Card**

You can use this card to help inform your doctors and nurses about all the medications that you take

This information is not a replacement for speaking with your healthcare team.

Sincerely,  
Genentech

# Understanding **YOUR DISEASE**



Being diagnosed with cancer can be overwhelming. It's okay if you don't know where to begin. Start by learning about hepatocellular carcinoma, or HCC, a type of liver cancer.

## Liver cancer is a **COMMON TYPE** of cancer

- 6th most common type of cancer in the world
- 13th most common in the United States

## **MOST LIVER CANCERS ARE HCC**

- 90% of liver cancers in the world are HCC
- 65% of cases in the United States are HCC

### Symptoms of HCC may include:

- |   |  |   |  |
|---|--|---|--|
|  Pain or swelling<br>in the belly |  Easy bruising<br>or bleeding |  Yellow skin or eyes<br>(jaundice) |  Unexplained<br>weight loss |
|---|--|---|--|

### How is HCC treated?

Some cases of HCC can be removed with surgery. Other cases cannot, and they can only be treated with medicine. Most medicines for HCC are systemic therapies. This means the medicine goes in your blood and travels all over your body. This type of treatment can be a pill, an injection, or an intravenous, or "IV," infusion. An infusion is a type of injection that's given over a period of time in a doctor's office or an infusion center.

### Systemic therapies for HCC

Beyond chemotherapy, there have been many medicines researched and approved to treat HCC in the last few years. Some treatments even combine medicines. Here is an overview of the types of medicine available:



#### **Targeted drug therapy**

- A commonly used type of advanced liver cancer treatment
- Some work to cut off blood and nutrients from reaching the tumor
- Some use other targets to help keep the cancer from growing and spreading



#### **Immunotherapy**

- Helps the immune system see cancer cells
- Designed to help find and fight the cancer

**Ask your doctor what types of treatment could be right for you**

Please see additional information on the following page.

## Should HCC be treated right away?

Once HCC is diagnosed, it's important to talk to your doctor right away about treatment options. If it isn't treated, the cancer can spread to other parts of the body.

### Some additional resources

**American Liver Foundation**

**[www.LiverFoundation.org](http://www.LiverFoundation.org)**  
1-800-465-4837

**Blue Faery, The Adrienne Wilson Liver Cancer Association**

**[www.BlueFaery.org](http://www.BlueFaery.org)**  
1-818-636-5624

**Global Liver Institute (GLI)**

**[www.GlobalLiver.org](http://www.GlobalLiver.org)**

**National Comprehensive Cancer Network (NCCN)**

**[www.NCCN.org/patients](http://www.NCCN.org/patients)**

Please be aware that links to third-party sites are provided for convenience only. Genentech is not responsible for the accuracy of the information contained on a third-party site and does not recommend or endorse the content provided on the site. Your use of third-party websites is at your own risk and subject to the terms and conditions of use for such sites.

## What else can I do?



**Consider changes to your lifestyle, habits, diet, and activity level.** Changes like these may help make you feel better during your cancer treatment. Ask your healthcare team what changes are right for you.

You can find resources in the Information and Support sheet included in this packet.

This is not intended to replace the advice of your healthcare team.

**Genentech**

*A Member of the Roche Group*

M-US-00005994(v1.0)

# Where to find more **INFORMATION AND SUPPORT**

There are lots of resources available to help people with liver cancer. Here is a list of some of the resources that may be available.



## **Cancer organizations**

These organizations are devoted to helping fight cancer through research, early detection, education, and treatment programs nationwide.

### **The American Cancer Society**

**[www.cancer.org](http://www.cancer.org)**

1-800-227-2345

The American Cancer Society's mission is to save lives, celebrate lives, and lead the fight for a world without cancer.

### **CancerCare**

**[www.cancercare.org](http://www.cancercare.org)**

1-800-813-4673

CancerCare provides telephone, online and face-to-face counseling, support groups, education, publications and financial and co-payment assistance.

### **Cancer.Net**

**[www.cancer.net](http://www.cancer.net)**

1-888-651-3038

Trusted, compassionate information for people with cancer and their families and caregivers, from the American Society of Clinical Oncology (ASCO), the voice of the world's cancer physicians and oncology professionals.

### **National Cancer Institute (NCI)**

**[www.cancer.gov](http://www.cancer.gov)**

1-800-422-6237

NCI's mission is to lead, conduct, and support cancer research across the nation to advance scientific knowledge and help all people live longer, healthier lives.

### **National Comprehensive Cancer Network (NCCN)**

**[www.nccn.org](http://www.nccn.org)**

1-215-690-0300

NCCN's mission is to improve and facilitate quality, effective, efficient, and accessible cancer care so patients can live better lives.

**Please see additional information on the following page.**



## Liver cancer organizations

These organizations are dedicated to helping those afflicted by liver cancer through a variety of resources, support groups, and outreach programs.

### **American Liver Foundation (ALF)**

**[www.LiverFoundation.org](http://www.LiverFoundation.org)**

1-800-465-4837

ALF's mission is to promote education, advocacy, support services and research for the prevention, treatment and cure of liver disease.

### **Blue Faery, The Adrienne Wilson Liver Cancer Association**

**[www.BlueFaery.org](http://www.BlueFaery.org)**

1-818-636-5624

Blue Faery's mission is to prevent, treat and cure primary liver cancer, specifically Hepatocellular Carcinoma, through research, education and advocacy.

### **Global Liver Institute**

**[www.globalliver.org](http://www.globalliver.org)**

Global Liver Institute wants to bring innovation and collaboration to the fight against liver diseases. We hope to address the gaps in attention, education, support, and research around improving liver health.



## Support organizations

These organizations are dedicated to providing care to cancer patients and their caregivers through social and emotional support efforts, educational programs, and wellness initiatives.

### **Cancer Support Community**

**[www.cancersupportcommunity.org](http://www.cancersupportcommunity.org)**

1-888-793-9355

Global network offering quality cancer support to millions of people touched by cancer, available online and at community-based centers and hospitals.

### **Patient Advocate Foundation**

**[www.patientadvocate.org](http://www.patientadvocate.org)**

1-800-532-5274

Patient Advocate Foundation (PAF) is a national 501 (c)(3) non-profit organization which provides case management services and financial aid to Americans with chronic, life threatening and debilitating illnesses.

### **Family Reach**

**[www.familyreach.org](http://www.familyreach.org)**

Boston office: 1-857-233-2764

New Jersey office: 1-973-394-1411

Family Reach is a national 501 (c)(3) organization dedicated to eradicating the financial barriers that accompany a cancer diagnosis.

Please be aware that links to third-party sites are provided for convenience only. Genentech is not responsible for the accuracy of the information contained on a third-party site and does not recommend or endorse the content provided on the site. Your use of third-party websites is at your own risk and subject to the terms and conditions of use for such sites.

**Genentech**

*A Member of the Roche Group*

M-US-00005994(v1.0)

# What to know about **TECENTRIQ + Avastin®** (bevacizumab)

## About TECENTRIQ

TECENTRIQ is a type of immunotherapy. That means it helps the immune system see cancer cells. It's designed to help find and fight the cancer. TECENTRIQ can cause your immune system to attack normal organs and tissues and can affect the way they work. These problems can sometimes become serious or life threatening and can lead to death. Your healthcare team will watch for signs of these problems. TECENTRIQ is **not chemotherapy**.

## TECENTRIQ + Avastin: proven results

Here are the results of a clinical study that compared TECENTRIQ + Avastin to sorafenib. The study included 501 patients with advanced hepatocellular carcinoma (HCC) or tumors that could not be removed by surgery.

### More patients **LIVED LONGER**

The study found patients taking TECENTRIQ + Avastin were 42% less likely to have died, compared to sorafenib.

### **LOWER CHANCE** of tumors growing or spreading

Patients taking TECENTRIQ + Avastin were 41% less likely to have their cancer get worse, compared to sorafenib.

## How is TECENTRIQ + Avastin given?

Your healthcare team will give TECENTRIQ + Avastin with an intravenous or "IV" line. This means a bag of each medicine is given in your vein, one after the other. TECENTRIQ + Avastin is given **every 3 weeks**. Your healthcare team will also decide how long you may be on treatment. If your disease gets worse or treatments can't be tolerated, your healthcare team may stop infusions.

### First infusion

The first **TECENTRIQ** infusion is given over **60 minutes**.



TECENTRIQ



60 minutes

### Following infusions

Depending on your response to the first infusion, later infusions of **TECENTRIQ** may take **30 minutes**.



TECENTRIQ



30 minutes

**More time will be needed for your other medications. Ask your healthcare team how long the entire process will take.**

## What is TECENTRIQ?

TECENTRIQ is a prescription medicine used to treat adults with:

**A type of liver cancer called hepatocellular carcinoma (HCC).**

- **TECENTRIQ may be used with the medicine bevacizumab when your liver cancer:**

- has spread or cannot be removed by surgery, **and**
- you have not received other medicines by mouth or injection through your vein (IV) to treat your cancer.

It is not known if TECENTRIQ is safe and effective in children.

## Important Safety Information

**What is the most important information about TECENTRIQ?**

TECENTRIQ can cause your immune system to attack normal organs and tissues and can affect the way they work. These problems can sometimes become serious or life threatening and can lead to death.

**Please see additional Important Safety Information on the following page and accompanying full Prescribing Information.**

**TECENTRIQ®**  
atezolizumab 840 mg / 1200 mg  
INJECTION FOR IV USE

## Important Safety Information (Continued)

**Call or see your healthcare provider right away if you get any symptoms of the following problems or these symptoms get worse.**

**TECENTRIQ can cause serious side effects, including:**

- **Lung problems (pneumonitis)**—signs and symptoms of pneumonitis may include new or worsening cough, shortness of breath, and chest pain
- **Liver problems (hepatitis)**—signs and symptoms of hepatitis may include yellowing of your skin or the whites of your eyes, severe nausea or vomiting, pain on the right side of your stomach area (abdomen), drowsiness, dark urine (tea colored), bleeding or bruising more easily than normal, and feeling less hungry than usual
- **Intestinal problems (colitis)**—signs and symptoms of colitis may include diarrhea (loose stools) or more bowel movements than usual, blood or mucus in your stools or dark, tarry, sticky stools, and severe stomach area (abdomen) pain or tenderness
- **Hormone gland problems (especially the thyroid, adrenal glands, pancreas, and pituitary)**—signs and symptoms that your hormone glands are not working properly may include headaches that will not go away or unusual headaches, extreme tiredness, weight gain or weight loss, dizziness or fainting, feeling more hungry or thirsty than usual, hair loss, changes in mood or behavior (such as decreased sex drive, irritability, or forgetfulness), feeling cold, constipation, your voice gets deeper, urinating more often than usual, nausea or vomiting, and stomach area (abdomen) pain
- **Problems in other organs**—signs and symptoms may include severe muscle weakness, numbness or tingling in hands or feet, confusion, blurry vision, double vision, or other vision problems, changes in mood or behavior, extreme sensitivity to light, neck stiffness, eye pain or redness, skin blisters or peeling, chest pain, irregular heartbeat, shortness of breath, or swelling of the ankles
- **Severe infections**—signs and symptoms of infection may include fever, cough, flu-like symptoms, pain when urinating, and frequent urination or back pain
- **Severe infusion reactions**—signs and symptoms of infusion reactions may include chills or shaking, itching or rash, flushing, shortness of breath or wheezing, swelling of your face or lips, dizziness, fever, feeling like passing out, and back or neck pain

**Getting medical treatment right away may help keep these problems from becoming more serious.** Your healthcare provider may treat you with corticosteroid or hormone replacement

medicines. Your healthcare provider may delay or completely stop treatment with TECENTRIQ if you have severe side effects.

**Before you receive TECENTRIQ, tell your healthcare provider about all of your medical conditions, including if you:**

- have immune system problems (such as Crohn's disease, ulcerative colitis, or lupus); have had an organ transplant; have lung or breathing problems; have liver problems; have a condition that affects your nervous system (such as myasthenia gravis or Guillain-Barre syndrome); or are being treated for an infection
- are pregnant or plan to become pregnant. TECENTRIQ can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with TECENTRIQ. **Females who are able to become pregnant:**
  - your healthcare provider should do a pregnancy test before you start treatment with TECENTRIQ
  - you should use an effective method of birth control during your treatment and for at least 5 months after the last dose of TECENTRIQ
- are breastfeeding or plan to breastfeed. It is not known if TECENTRIQ passes into your breast milk. Do not breastfeed during treatment and for at least 5 months after the last dose of TECENTRIQ

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**The most common side effects of TECENTRIQ when used in hepatocellular carcinoma (HCC) with bevacizumab include:**

- high blood pressure
- feeling tired or weak
- too much protein in the urine

TECENTRIQ may cause fertility problems in females, which may affect the ability to have children. Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of TECENTRIQ. Ask your healthcare provider or pharmacist for more information about the benefits and side effects of TECENTRIQ.

You may report side effects to the FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch). You may also report side effects to Genentech at 1-888-835-2555.

**Please see accompanying full Prescribing Information for additional Important Safety Information.**

# Options that can help you **AFFORD TREATMENT**

## Genentech Access Solutions has options that may help you afford your prescribed Genentech medicine

Genentech Access Solutions helps you understand your insurance coverage and options for financial assistance. This can help you get the Genentech medicine your doctor prescribed.



Help you understand the details of your coverage, such as:

- If your health insurance will pay for your Genentech medicine
- How much your co-pay will cost for your medicine



Refer you to financial assistance options:

- Genentech Patient Foundation
- Genentech co-pay programs
- Independent co-pay assistance foundations



Visit **Genentech-Access.com** to learn how Genentech Access Solutions may be able to help you

## Genentech Patient Foundation

**If you don't have insurance** or are worried about paying for medicine, the Genentech Patient Foundation might be able to get you free medicine if you are eligible.

### Genentech Patient Foundation eligibility criteria

#### UNINSURED PATIENTS

With incomes under \$150,000

OR

#### INSURED PATIENTS WITHOUT COVERAGE

for a Genentech medicine  
With incomes under \$150,000

OR

#### INSURED PATIENTS WITH COVERAGE

for a Genentech medicine

- With unaffordable out-of-pocket costs
- Who have used all forms of available financial assistance
- With household size and income within certain guidelines, available at **GenentechPatientFoundation.com**

Please see additional information on the following page.

## Genentech co-pay programs

Genentech co-pay programs help people with **commercial health insurance**. This might be a plan you get through your employer or one you purchased through a Health Insurance Marketplace like HealthCare.gov. To qualify, you must also meet other criteria.

You pay as little as \$5 for Genentech medications, if you qualify for the Genentech Oncology Co-pay Card. The program covers the rest, up to a \$25,000 yearly limit per Genentech medication.\*

<b>NO</b> INCOME REQUIREMENTS	<b>PAY AS LITTLE AS</b> <b>\$5</b> PER CO-PAY	<b>\$25k</b> ANNUAL BENEFIT LIMIT PER GENENTECH PRODUCT
----------------------------------	---	---

\*Eligibility criteria apply. Not valid for patients using federal or state government programs to pay for their medications. Patient must be taking the Genentech medication for an FDA-approved indication. See full terms and conditions at [CoPayAssistanceNow.com](https://www.CoPayAssistanceNow.com).



Learn more about this program at [CoPayAssistanceNow.com](https://www.CoPayAssistanceNow.com)

## Independent co-pay assistance foundations

Independent co-pay assistance foundations are charitable organizations that give financial assistance for medicines. They help patients with public or commercial health insurance. You do not need to be enrolled in Genentech Access Solutions to be referred to an independent co-pay assistance foundation.



You can visit [Genentech-Access.com](https://www.Genentech-Access.com) to view a list of independent co-pay assistance foundations

OR



Call Genentech Access Solutions at **(888) 249-4918** to get help



### Still have questions?

Call the Genentech Patient Resource Center at **(877) GENENTECH (436-3683)** Monday-Friday, 6AM-5PM PST with questions or to get started.

# Learn about **POTENTIAL SIDE EFFECTS**

As you get started with TECENTRIQ + Avastin® (bevacizumab), you may feel the signs and symptoms of the side effects listed below. **Call or see your healthcare provider right away if you feel any symptoms of the following side effects or if these symptoms get worse.** You can find more information about these side effects on the back of this sheet.



## Symptoms of side effects may include:



### Lung problems (pneumonitis)

- New or worsening cough
- Shortness of breath
- Chest pain



### Liver problems (hepatitis)

- Yellowing of your skin or the whites of your eyes
- Severe nausea or vomiting
- Pain on the right side of your stomach area (abdomen)
- Drowsiness
- Dark urine (tea colored)
- Bleeding or bruising more easily than normal
- Feeling less hungry than usual



### Intestinal problems (colitis)

- Diarrhea (loose stools) or more bowel movements than usual
- Blood or mucus in your stools or dark, tarry, sticky stools
- Severe stomach area (abdomen) pain or tenderness



### Hormone gland problems (especially the thyroid, adrenal glands, pancreas, and pituitary)

- Headaches that will not go away or unusual headaches
- Extreme tiredness
- Weight gain or weight loss
- Dizziness or fainting
- Feeling more hungry or thirsty than usual
- Hair loss
- Changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness
- Feeling cold
- Constipation
- Your voice gets deeper
- Urinating more often than usual
- Nausea or vomiting
- Stomach area (abdomen) pain



### Problems in other organs

- Severe muscle weakness
- Numbness or tingling in hands or feet
- Confusion
- Blurry vision, double vision, or other vision problems
- Changes in mood or behavior
- Extreme sensitivity to light
- Neck stiffness
- Eye pain or redness
- Skin blisters or peeling
- Chest pain, irregular heartbeat, shortness of breath, or swelling of the ankles



### Severe infections

- Fever
- Cough
- Flu-like symptoms
- Pain when urinating, frequent urination, or back pain



### Severe infusion reactions

- Chills or shaking
- Itching or rash
- Flushing
- Shortness of breath or wheezing
- Swelling of your face or lips
- Dizziness
- Fever
- Feeling like passing out
- Back or neck pain

## Report any changes you notice



Your healthcare team will likely keep a record of your side effects. Be sure to connect with them to keep track of symptoms and how you're feeling.

This information is not intended to be medical advice or replace the advice of your healthcare team.

Please see Important Safety Information on the following page and accompanying full Prescribing Information.

 **TECENTRIQ®**  
atezolizumab 840 mg / 1200 mg  
INJECTION FOR IV USE

## What is TECENTRIQ?

TECENTRIQ is a prescription medicine used to treat adults with:

### A type of liver cancer called hepatocellular carcinoma (HCC).

- **TECENTRIQ may be used with the medicine bevacizumab when your liver cancer:**
  - has spread or cannot be removed by surgery, and
  - you have not received other medicines by mouth or injection through your vein (IV) to treat your cancer.

It is not known if TECENTRIQ is safe and effective in children.

## Important Safety Information

### What is the most important information about TECENTRIQ?

TECENTRIQ can cause your immune system to attack normal organs and tissues and can affect the way they work. These problems can sometimes become serious or life threatening and can lead to death.

**Call or see your healthcare provider right away if you get any symptoms of the following problems or these symptoms get worse.**

**TECENTRIQ can cause serious side effects, including:**

- **Lung problems (pneumonitis)**—signs and symptoms of pneumonitis may include new or worsening cough, shortness of breath, and chest pain
- **Liver problems (hepatitis)**—signs and symptoms of hepatitis may include yellowing of your skin or the whites of your eyes, severe nausea or vomiting, pain on the right side of your stomach area (abdomen), drowsiness, dark urine (tea colored), bleeding or bruising more easily than normal, and feeling less hungry than usual
- **Intestinal problems (colitis)**—signs and symptoms of colitis may include diarrhea (loose stools) or more bowel movements than usual, blood or mucus in your stools or dark, tarry, sticky stools, and severe stomach area (abdomen) pain or tenderness
- **Hormone gland problems (especially the thyroid, adrenal glands, pancreas, and pituitary)**—signs and symptoms that your hormone glands are not working properly may include headaches that will not go away or unusual headaches, extreme tiredness, weight gain or weight loss, dizziness or fainting, feeling more hungry or thirsty than usual, hair loss, changes in mood or behavior (such as decreased sex drive, irritability, or forgetfulness), feeling cold, constipation, your voice gets deeper, urinating more often than usual, nausea or vomiting, and stomach area (abdomen) pain
- **Problems in other organs**—signs and symptoms may include severe muscle weakness, numbness or tingling in hands or feet, confusion, blurry vision, double vision, or other vision problems, changes in mood or behavior, extreme sensitivity to light, neck stiffness, eye pain or redness, skin blisters or peeling, chest pain, irregular heartbeat, shortness of breath, or swelling of the ankles
- **Severe infections**—signs and symptoms of infection may include fever, cough, flu-like symptoms, pain when urinating, and frequent urination or back pain

- **Severe infusion reactions**—signs and symptoms of infusion reactions may include chills or shaking, itching or rash, flushing, shortness of breath or wheezing, swelling of your face or lips, dizziness, fever, feeling like passing out, and back or neck pain

**Getting medical treatment right away may help keep these problems from becoming more serious.** Your healthcare provider may treat you with corticosteroid or hormone replacement medicines. Your healthcare provider may delay or completely stop treatment with TECENTRIQ if you have severe side effects.

**Before you receive TECENTRIQ, tell your healthcare provider about all of your medical conditions, including if you:**

- have immune system problems (such as Crohn's disease, ulcerative colitis, or lupus); have had an organ transplant; have lung or breathing problems; have liver problems; have a condition that affects your nervous system (such as myasthenia gravis or Guillain-Barre syndrome); or are being treated for an infection
- are pregnant or plan to become pregnant. TECENTRIQ can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with TECENTRIQ. **Females who are able to become pregnant:**
  - your healthcare provider should do a pregnancy test before you start treatment with TECENTRIQ
  - you should use an effective method of birth control during your treatment and for at least 5 months after the last dose of TECENTRIQ
- are breastfeeding or plan to breastfeed. It is not known if TECENTRIQ passes into your breast milk. Do not breastfeed during treatment and for at least 5 months after the last dose of TECENTRIQ

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**The most common side effects of TECENTRIQ when used in hepatocellular carcinoma (HCC) with bevacizumab include:**

- high blood pressure
- feeling tired or weak
- too much protein in the urine

TECENTRIQ may cause fertility problems in females, which may affect the ability to have children. Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of TECENTRIQ. Ask your healthcare provider or pharmacist for more information about the benefits and side effects of TECENTRIQ.

You may report side effects to the FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch). You may also report side effects to Genentech at 1-888-835-2555.

**Please see accompanying full Prescribing Information for additional Important Safety Information.**

**Genentech**

A Member of the Roche Group

M-US-00005994(v1.0)

 **TECENTRIQ**<sup>®</sup>  
atezolizumab 840 mg / 1200 mg  
INJECTION FOR IV USE

# Ways to help **MANAGE SIDE EFFECTS**

As you get started, it's helpful to know what to expect from treatment. TECENTRIQ + Avastin® (bevacizumab) may cause side effects. **Talk to your healthcare team about any side effects you have.**



## Feeling tired or weak

- **Plan time to rest.** Short naps may help. But don't sleep too much during the day
- **Balance rest and activity.** Short walks and light exercise may help. It's also important to relax when possible
- **Meet with a dietitian.** He or she can help you learn about foods and drinks that can boost your energy



## Diarrhea

- **Drink fluids.** Diarrhea can make you dehydrated, so be sure to drink more liquids
- **Eat small meals.** Break your mealtime into 6 to 8 smaller meals
- **Choose different foods.** Try the "BRAT" diet. Eat Bananas, Rice, Applesauce, and Toast in small amounts



## Decreased appetite

- **Track what you eat.** Share your notes with your healthcare team. You may find that certain foods or mealtimes work better than others
- **Eat several light meals.** Replace regular mealtimes with 5 or 6 smaller meals
- **Stay active.** Even a short walk can help increase your appetite



## Build a healthy diet

- **Choose foods that work for you.** Certain foods or mealtimes may work better for you
- **Drink fluids.** You should take in 8 to 12 cups of fluids every day
- **Meet with a dietitian.** He or she can suggest changes to lessen side effects and help you maintain a healthy weight



## Balance rest with exercise

- **Choose activities you like.** For example: If you like bicycling, look for ways to add it into your daily life
- **Make small changes.** Even a short walk can make a difference
- **Try to do as much as before.** If you already have an exercise routine, see if you can continue it

## What is TECENTRIQ?

TECENTRIQ is a prescription medicine used to treat adults with a **type of liver cancer called hepatocellular carcinoma (HCC)**. **TECENTRIQ may be used with the medicine bevacizumab when your liver cancer** has spread or cannot be removed by surgery, **and** you have not received other medicines by mouth or injection through your vein (IV) to treat your cancer.

It is not known if TECENTRIQ is safe and effective in children.

Please see additional Important Safety Information on the following page and accompanying full [Prescribing Information](#).

 **TECENTRIQ**<sup>®</sup>  
atezolizumab 840 mg / 1200 mg  
INJECTION FOR IV USE

## Important Safety Information

**What is the most important information about TECENTRIQ?** TECENTRIQ can cause your immune system to attack normal organs and tissues and can affect the way they work. These problems can sometimes become serious or life threatening and can lead to death.

**Call or see your healthcare provider right away if you get any symptoms of the following problems or these symptoms get worse.**

**TECENTRIQ can cause serious side effects, including** lung problems (pneumonitis), liver problems (hepatitis), intestinal problems (colitis), hormone gland problems (especially the thyroid, adrenal glands, pancreas, and pituitary), problems in other organs, severe infections, and severe infusion reactions.

**Getting medical treatment right away may help keep these problems from becoming more serious.** Your healthcare provider may delay or completely stop treatment with TECENTRIQ if you have severe side effects.

**Before you receive TECENTRIQ, tell your healthcare provider about all of your medical conditions, including if you:**

- have immune system problems; have had an organ transplant; have lung or breathing problems; have liver problems; have a condition that affects your nervous system; or are being treated for an infection
- are pregnant or plan to become pregnant. TECENTRIQ can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with TECENTRIQ.

**Females who are able to become pregnant:**

- your healthcare provider should do a pregnancy test before you start treatment with TECENTRIQ
  - you should use an effective method of birth control during your treatment and for at least 5 months after the last dose of TECENTRIQ
- are breastfeeding or plan to breastfeed. It is not known if TECENTRIQ passes into your breast milk. Do not breastfeed during treatment and for at least 5 months after the last dose of TECENTRIQ

**Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.**

**The most common side effects of TECENTRIQ when used in hepatocellular carcinoma (HCC) with bevacizumab include:**

- high blood pressure
- feeling tired or weak
- too much protein in the urine

TECENTRIQ may cause fertility problems in females, which may affect the ability to have children. Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of TECENTRIQ. Ask your healthcare provider or pharmacist for more information about the benefits and side effects of TECENTRIQ.

You may report side effects to the FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch). You may also report side effects to Genentech at 1-888-835-2555.

**Please see accompanying full Prescribing Information for additional Important Safety Information.**

### Other possible side effects & tips



Treatment may cause other side effects. **Find a list in the accompanying Medication Guide.**



Talk to your healthcare team about your side effects and how to manage them. You can also find tips for managing side effects of cancer treatment at: **[Cancer.gov/about-cancer/treatment/side-effects](https://www.cancer.gov/about-cancer/treatment/side-effects)**

## MY MEDICATION CARD



My name \_\_\_\_\_

(   )   -

Phone number

Oncologist name \_\_\_\_\_

(   )   -

Phone number

Hepatologist name \_\_\_\_\_

(   )   -

Phone number

Primary care doctor name \_\_\_\_\_

(   )   -

Phone number

Emergency contact \_\_\_\_\_

(   )   -

Phone number

I use TECENTRIQ + Avastin® (bevacizumab)  
to treat my: \_\_\_\_\_



Here are other medications I take: \_\_\_\_\_

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

**Note to healthcare team:** If you have been given this card by a patient who is on TECENTRIQ treatment, this information is not intended to replace your medical judgment.

M-US-00005994(v1.0)