



[pronunciation zep zel' kah]

Get ready for the next step

Your guide for treatment
on ZEPZELCA™ (lurbinectedin)



What is ZEPZELCA?

ZEPZELCA is a prescription medicine used to treat adults with a kind of lung cancer called small cell lung cancer (SCLC).

ZEPZELCA may be used when your lung cancer:

- has spread to other parts of the body (metastatic), **and**
- you have received treatment with chemotherapy that contains platinum, and it did not work or is no longer working.

ZEPZELCA is approved based on response rate and how long the response lasted. Additional studies will further evaluate the benefit of ZEPZELCA for this use.

IMPORTANT SAFETY INFORMATION

ZEPZELCA can cause serious side effects, including low blood cell counts and liver problems. Before receiving ZEPZELCA, tell your doctor if you are pregnant or plan to become pregnant. ZEPZELCA can harm your unborn baby. These are not all of the possible side effects of ZEPZELCA. See additional Important Safety Information on pages 8, 9, and 11.

Please see accompanying full Prescribing Information including Patient Information, and discuss with your doctor.

Is this guide for you?

This guide is for you if:

- You have been prescribed ZEPZELCA™ (lurbinectedin)
- Your loved one has been prescribed ZEPZELCA
- You want to know more about ZEPZELCA

This guide can help you:

- Know what to expect from treatment with ZEPZELCA
- Get the most from your treatment

Words in **maroon** in this guide are defined on page 14.

Table of Contents

04 | THE BASICS

- Small cell lung cancer and its treatment

06 | WHAT ZEPZELCA DOES

- What is ZEPZELCA?
- How does ZEPZELCA work?

08 | SIDE EFFECTS

- What are the possible side effects of ZEPZELCA?

10 | TAKING ZEPZELCA

- How will I receive ZEPZELCA?
- What should I tell my doctor before receiving ZEPZELCA?

12 | SUPPORT


- How do I get access and support for ZEPZELCA?
- Support groups and resources

14 | GLOSSARY/NOTES

This guide does not replace your doctor's advice.
Please ask them any questions you have about treatment.

Please see accompanying full Prescribing Information
including Patient Information, and discuss with your doctor.

The basics: Small cell lung cancer



You've already been through treatment for small cell lung cancer. You probably know a good bit about it already. Here are some reminders and maybe some new terms that can help you understand treatment with ZEPZELCA™ (lurbinectedin).

About SCLC

What is small cell lung cancer?

Cancer starts when abnormal cells in the body grow out of control. Lung cancer starts in the lungs.

There are 2 types of lung cancer. One is called non–small cell lung cancer. The other is called small cell lung cancer, or SCLC.

SCLC tends to grow and spread quickly. That's why it tends to respond to **chemotherapy** and radiation therapy. For the same reason, SCLC comes back for most people.

What is **metastatic** lung cancer?

Cancer that has spread from the lungs, where it started, to other parts of the body.

What is progression?

Disease that has spread and grown.

Please see accompanying full Prescribing Information including Patient Information, and discuss with your doctor.

Treatment for SCLC

First-line chemotherapies for SCLC

SCLC is usually first treated with **chemotherapy** combinations.

The combinations used most often are:

- Cisplatin + etoposide or irinotecan
- Carboplatin + etoposide or irinotecan

SCLC often responds to first-line chemotherapy. But it usually comes back sooner or later.



When SCLC comes back

There are still treatment options when the first treatment did not work or is no longer working.

ZEPZELCA is used by itself in adults:

- With SCLC that has spread and
- Who have already tried cisplatin or carboplatin

It can be hard to hear that your SCLC has come back.
But ZEPZELCA can help you keep fighting.

What is ZEPZELCA™ (lurbinectedin)?

ZEPZELCA is a prescription medicine used to treat adults with a kind of lung cancer called small cell lung cancer (SCLC).

ZEPZELCA may be used when your lung cancer:

- has spread to other parts of the body (**metastatic**), and
- you have received treatment with **chemotherapy** that contains platinum, and it did not work or is no longer working

ZEPZELCA is approved based on **response** rate and how long the response lasted. Additional studies will further evaluate the benefit of ZEPZELCA for this use.

How does ZEPZELCA work?

ZEPZELCA works in the body to help fight metastatic SCLC by:

1


Binding to **DNA**

2

Interfering with cancer cell function

3

Causing cancer cells to die



What are the possible side effects of ZEPZELCA™ (lurbinectedin)?

ZEPZELCA can cause serious side effects, including:

- **Low blood cell counts.** Low blood counts including low neutrophil counts (neutropenia) and low platelet counts (thrombocytopenia) are common with ZEPZELCA, and can also be severe. Some people with low white blood cell counts may get fever, or an infection throughout the body (sepsis), that can cause death. Your healthcare provider should do blood tests before you receive each treatment with ZEPZELCA to check your blood cell counts.

Tell your healthcare provider right away if you develop:

- fever or any other signs of infection
- tiredness
- unusual bruising or bleeding
- pale colored skin

- **Liver problems.** Increased liver function tests are common with ZEPZELCA, and can also be severe. Your healthcare provider should do blood tests to check your liver function before you start and during treatment with ZEPZELCA.

Tell your healthcare provider right away if you develop symptoms of liver problems including:

- loss of appetite
- nausea or vomiting
- pain on the right side of your stomach area (abdomen)

Your healthcare provider may temporarily stop treatment, lower your dose, or permanently stop ZEPZELCA if you develop low blood cell counts or liver problems during treatment with ZEPZELCA.

The most common side effects of ZEPZELCA include:


- tiredness
- low white and red blood cell counts
- increased kidney function blood test (creatinine)
- increased liver function blood tests
- increased blood sugar (glucose)
- nausea
- decreased appetite
- muscle and joint (musculoskeletal) pain
- low level of albumin in the blood
- constipation
- trouble breathing
- low levels of sodium and magnesium in the blood
- vomiting
- cough
- diarrhea

These are not all of the possible side effects of ZEPZELCA. **For complete safety information, please see accompanying full Prescribing Information including Patient Information, and discuss with your doctor.**

Call your doctor for medical advice about side effects.
You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call **1-800-FDA-1088**. You may also report side effects to Jazz Pharmaceuticals at **1-800-520-5568**.

How will I receive ZEPZELCA™ (lurbinectedin)?

ZEPZELCA is usually given **over 1 hour every 3 weeks**:



Before each treatment with ZEPZELCA:

- You may receive medicines to help prevent nausea and vomiting or make it less severe

During your treatment with ZEPZELCA:

- Your healthcare provider may do certain tests to check you for side effects, and to see how well you respond to the treatment

How long will I be on treatment?

- Your doctor will decide how long you will continue treatment with ZEPZELCA

With ZEPZELCA, you have time between treatments to focus on what you want.

What should I tell my doctor before receiving ZEPZELCA?

Tell your healthcare provider about all of your medical conditions, including if you:

- have liver or kidney problems
- are pregnant or plan to become pregnant. ZEPZELCA can harm your unborn baby

Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with ZEPZELCA
- You should use effective birth control (contraception) during treatment with and for 6 months after your final dose of ZEPZELCA
- Tell your healthcare provider right away if you become pregnant or think that you are pregnant during treatment with ZEPZELCA

Males with female partners who are able to become pregnant should use effective birth control during treatment with and for 4 months after your final dose of ZEPZELCA.

- are breastfeeding or plan to breastfeed. It is not known if ZEPZELCA passes into your breastmilk. Do not breastfeed during treatment with ZEPZELCA and for 2 weeks after your final dose of ZEPZELCA. Talk to your healthcare provider about the best way to feed your baby during treatment with ZEPZELCA

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Certain other medicines may affect how ZEPZELCA works.

What should I avoid while using ZEPZELCA?

- Avoid eating or drinking grapefruit, or products that contain grapefruit juice, during treatment with ZEPZELCA

Please see accompanying full Prescribing Information including Patient Information, and discuss with your doctor.

How do I get access and support for ZEPZELCA™ (lurbinectedin)?



JazzCares is committed to helping you get access to your ZEPZELCA medication



To help you get access to ZEPZELCA, JazzCares specialists are available to assist you with

- Understanding your insurance coverage
- Other cost and coverage questions
- Referrals to other financial assistance



JazzCares is committed to helping you reduce your out-of-pocket costs for ZEPZELCA

- Savings Card—eligible, commercially insured patients can pay as little as \$10 for their ZEPZELCA medication, subject to an annual maximum



Free drug program, if you are eligible

- Financial and other eligibility criteria may apply

Learn more about JazzCares support offerings by calling 1-833-533-JAZZ (5299) Monday–Friday between 8 AM and 8 PM ET or visiting [JazzCares.com](https://www.jazzcares.com)

Insurance coverage and plans may vary. The JazzCares program at Jazz Pharmaceuticals provides general information only and is not a guarantee of any coverage or reimbursement outcome. All treatment decisions rest solely with the treating physician or qualified healthcare professional. Jazz Pharmaceuticals reserves the right to terminate or modify this program at any time with or without notice. Other terms and conditions apply.

Support groups and resources

Having cancer isn't easy. When it comes back, it can be even harder.

Remember what you can do:

- Gain knowledge to help you feel in control of your decisions
- Talk with others who understand what you're going through

A number of helpful resources and support groups are here for you. These organizations are independent of Jazz Pharmaceuticals.

American Cancer Society® Everything from emotional support to the latest cancer information. Plus information about support programs in your area. www.cancer.org

CancerCare® Free, professional support services for people affected by lung cancer. www.cancercare.org

Cancer Support Community Online cancer support, including discussion boards, counselors, blog platforms, and more. www.cancersupportcommunity.org

GO₂ Foundation for Lung Cancer Dedicated to saving, extending, and improving the lives of people with lung cancer. www.go2foundation.org

Lungevity Information, resources, and ways to connect with patients, caregivers, and medical experts. www.lungevity.org

Lung Cancer Research Foundation® Educational materials and a lung cancer support line. www.lungcancerresearchfoundation.org

Lung Cancer Foundation of America A resource for patients seeking answers, hope, and access to updated treatment information. www.lcfamerica.org

All product names and brands are property of their respective owners.

Please see accompanying full Prescribing Information including Patient Information, and discuss with your doctor.

Glossary of terms

Chemotherapy: Treatment with drugs that kill cancer cells. They may be injected into a vein or taken by mouth. These drugs travel through the bloodstream and reach most parts of the body.

DNA: The genetic blueprint found in the center of each cell. DNA holds the instructions for cell growth, division, and function. DNA stands for deoxyribonucleic acid.

First-line treatment: The first therapy given after a cancer diagnosis.

Metastatic: Cancer that has spread from where it started to other structures or organs in the body.

Response: The outcome after treatment, or the reaction a patient has to a drug or any other therapy.

Add any words you want to remember:

Have questions for your doctor or nurse?

Use this space to write down any questions you have about treatment with ZEPZELCA™ (lurbinectedin). Remember to bring this with you to your next doctor’s appointment.

Key Contacts:

Name: _____

Number: _____

Email: _____

Name: _____

Number: _____

Email: _____

Get the most from your treatment with



Visit us at **ZEPZELCA.com**

Learn more about patient support and access
to ZEPZELCA at **JazzCares.com**

**Please see accompanying full Prescribing Information including
Patient Information, and discuss with your doctor.**



Jazz Pharmaceuticals[®]



ZEPZELCA is a trademark of PharmaMar, S.A. used by Jazz Pharmaceuticals under license.

©2021 Jazz Pharmaceuticals US-LUR-2000165 Rev0121