

# UKONIQ™

umbralisib 200 mg  
tablets

## GOT THIS

### YOUR GUIDE TO TREATMENT WITH UKONIQ

UKONIQ is a prescription medicine used to treat adults with:

- **Marginal zone lymphoma (MZL)** when the disease has come back or did not respond to treatment and who have received at least 1 certain type of prior treatment
- **Follicular lymphoma (FL)** when the disease has come back or did not respond to treatment and who have received at least 3 prior treatments

*Not an actual patient.*

It is not known if UKONIQ is safe and effective in children. UKONIQ was approved through a faster FDA review process based on the percentage of patients whose tumor size shrank or disappeared after treatment. There is an ongoing study to confirm the benefit of UKONIQ for these uses.

### IMPORTANT SAFETY INFORMATION

**UKONIQ can cause serious side effects, including:**

**Infections.** UKONIQ can cause serious infections that may lead to death. Tell your healthcare provider right away if you have fever, chills, weakness, flu-like symptoms, or any other signs of infection during treatment with UKONIQ.

**Please see Important Safety Information throughout and Important Facts about UKONIQ on pages 10-11.**

## Living with follicular lymphoma (FL)

### FL is the most common slow-growing non-Hodgkin's lymphoma

If you're living with FL, slow-growing cancer cells may be found in your lymph nodes. The cells often appear to be in rounded clumps that look like follicles, which is why this type of lymphoma is called "follicular."

#### What happens when FL comes back?

FL is a chronic disease that many people live with for a long time. However, every person responds to treatment differently. It is not uncommon for FL to come back, even years after treatment.

While FL grows slowly, symptoms can include:

- Enlarged or swollen lymph nodes in the neck, armpit, or groin
- Weight loss
- Fevers or night sweats
- Fatigue
- Susceptibility to infection



### IMPORTANT SAFETY INFORMATION (CONT'D)

**Low white blood cell count (neutropenia).** Neutropenia is common with UKONIQ treatment and can sometimes be serious. Your healthcare provider will check your blood counts regularly during treatment with UKONIQ. Tell your healthcare provider right away if you have a fever or any signs of infection during treatment with UKONIQ.

**Diarrhea or inflammation of your intestine (colitis).** Diarrhea is common during UKONIQ treatment and can sometimes be serious. Tell your healthcare provider right away if you have diarrhea that does not go away or worsening diarrhea, stool with mucus or blood, or if you have severe stomach area (abdominal) pain during treatment with UKONIQ. Drink plenty of fluids during treatment with UKONIQ to help prevent dehydration from diarrhea.

## Living with marginal zone lymphoma (MZL)

### MZL is the second most common slow-growing NHL

There are 3 types of MZL: extranodal, nodal, and splenic. Depending on the type of MZL you have, cancerous cells could be in the stomach or intestine, the spleen, in lymph nodes, or other parts of the body.

#### What happens when MZL comes back?

MZL is a chronic disease that many people live with for a long time. Sometimes MZL can be treated with antibiotics, or it can even go untreated. Some people may require chemotherapy and/or another type of infusion or oral therapy. Every person responds to treatment differently, which is why MZL may come back even years after treatment.

Symptoms are different depending on the type of MZL and can include:

- Enlarged or swollen lymph nodes
- Stomach pain or chronic indigestion
- Fatigue
- Enlarged spleen
- Anemia
- Low platelet counts
- Weight loss
- Fever

### Sometimes people do not experience symptoms when an indolent lymphoma comes back

It's important that you continue to be monitored by your doctor so they can look for signs and/or symptoms of your lymphoma.

Please see Important Safety Information throughout and Important Facts about UKONIQ on pages 10-11.

**UKONIQ**<sup>™</sup>  
umbralisib 200 mg  
tablets

# Get to know UKONIQ (YOU-KON-IK)

## What is UKONIQ?

**UKONIQ is a targeted therapy, not a chemotherapy.** UKONIQ is a prescription medicine used to treat adults with MZL when the disease has come back or did not respond to treatment and who have received at least 1 certain type of prior treatment and FL when the disease has come back or did not respond to treatment and who have received at least 3 prior treatments.

It is not known if UKONIQ is safe and effective in children. UKONIQ was approved through a faster FDA review process based on the percentage of patients whose tumor size shrank or disappeared after treatment. There is an ongoing study to confirm the benefit of UKONIQ for these uses.

## UKONIQ works differently

UKONIQ is a targeted therapy, not a chemotherapy, that blocks at least 2 different enzymes found in cancerous cells.



## IMPORTANT SAFETY INFORMATION (CONT'D)

**Liver problems.** Abnormal liver function blood test results are common and can sometimes be serious. Your healthcare provider will do blood tests before and during your treatment with UKONIQ to check for liver problems. Tell your healthcare provider right away if you have any of the following symptoms of liver problems:

- Yellowing of your skin or the white part of your eyes (jaundice)
- Dark or brown (tea colored) urine
- Pain in the upper right side of your stomach area (abdomen)
- Bleeding or bruising more easily than normal

# Starting UKONIQ

## Once-daily medication you can take at home



**800 mg taken orally  
(four 200 mg tablets)**



**Once daily at the same time**



**Take with food and swallow whole. Do not crush, break, cut, or chew the tablets**

## Here are some tips for remembering to take UKONIQ:

- **Keep UKONIQ in a safe place out of the reach of children**
- **Write yourself a reminder note**
- **Set an alarm for the same time each day**
- **Have someone remind you to take UKONIQ**



**Reminder:** If you miss a dose of UKONIQ, take it as soon as you remember on the same day. If it has been more than 12 hours, skip the missed dose and take your next dose on the next day at your usual time.

**Take UKONIQ exactly as your healthcare provider tells you. Your healthcare provider may tell you to decrease your dose, temporarily stop, or completely stop taking UKONIQ if you develop side effects. Do not change your dose or stop taking UKONIQ without talking to your healthcare provider first. If you vomit after taking a dose of UKONIQ, do not take another dose on that day. Take your next dose at your usual time.**

**Please see Important Safety Information throughout and Important Facts about UKONIQ on pages 10-11.**

**UKONIQ™**  
umbralisib 200 mg  
tablets

# The power of UKONIQ

**UKONIQ was tested in a clinical trial in adults with FL or MZL whose disease had come back or did not respond to treatment after 1 or 2 prior medicines**

**In the clinical trial that tested UKONIQ:**

**TUMORS SHRUNK OR COMPLETELY DISAPPEARED IN ALMOST HALF OF PATIENTS**



Tumors shrunk or completely disappeared in 50 patients (43%) with FL and in 34 patients (49%) with MZL. Tumors completely disappeared in 4 patients (3%) with FL and in 11 patients (16%) with MZL. Tumors shrunk in 46 patients (39%) with FL and in 23 patients (33%) with MZL. Half of the patients with FL who responded to UKONIQ had a response lasting more than 11 months and half who responded had a response lasting less than 11 months (called median duration of response). For patients with MZL, a median duration of response has not yet been determined because more than half of patients are still responding.

## IMPORTANT SAFETY INFORMATION (CONT'D)

**Severe skin reactions.** Rashes and other skin reactions are common with UKONIQ treatment and can sometimes be severe and lead to death. Tell your healthcare provider right away if you get a new or worsening skin rash or other signs of a severe skin reaction during treatment with UKONIQ, including:

- Painful sores or ulcers on your skin, lips, or in your mouth
- Rash with blisters or peeling skin
- Rash with itching
- Rash with fever

# What to expect when taking UKONIQ

**Talk to your doctor about how you're feeling before, during, and after treatment**

It's important to talk to your doctor or nurse right away about anything you may be feeling or experiencing.

**The most common side effects of UKONIQ include:**

- Changes in certain kidney function blood tests
- Tiredness
- Nausea
- Muscle or bone pain
- Low red blood cell count (anemia)
- Low platelet count
- Upper respiratory tract infection
- Vomiting
- Abdominal pain
- Decreased appetite

**Diarrhea is a common side effect of UKONIQ and can sometimes be serious. Guidelines to help manage diarrhea include:**

- Drink plenty of fluids during treatment with UKONIQ to help prevent dehydration from diarrhea
- Tell your doctor if you have diarrhea that does not go away or gets worse, stool with mucus or blood, or if you have severe stomach area (abdominal) pain during treatment with UKONIQ
- Keep track of your stools using a written tracker or by downloading a stool tracker app on your smartphone
- Visit [UKONIQ.com](https://www.ukoniq.com) to download a printable stool tracker or a digital version compatible with your smartphone



## Reminders before taking UKONIQ

- Let your doctor and nurses know about the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements
- Tell your doctor about all of your medical conditions, including if you: have an infection, have abdominal or intestinal problems, have liver problems, are allergic to FD&C Yellow No. 5 (tartrazine) or aspirin, are pregnant or plan to become pregnant, or are breastfeeding

**Please see Important Safety Information throughout and Important Facts about UKONIQ on pages 10-11.**

**UKONIQ™**  
umbralisib 200 mg tablets

## The power to stay informed

### These organizations will help support you through your treatment

**CancerCare**  
1-800-813-HOPE (1-800-813-4673)  
cancercares.org

**Cancer Support Community**  
1-888-793-WELL (1-888-793-9355)  
cancersupportcommunity.org

**The Leukemia & Lymphoma Society**  
1-800-955-4572  
lls.org

**Lymphoma Research Foundation**  
1-800-500-9976  
lymphoma.org

**National Coalition for Cancer Survivorship**  
1-877-NCCS-YES  
canceradvocacy.org

**Patient Empowerment Network**  
powerfulpatients.org

**Patient Power**  
patientpower.info

**Triage Cancer**  
1-424-258-4628  
trriagecancer.org

\*Please note that these are third-party organizations that are unaffiliated with TG Therapeutics. TG Therapeutics does not necessarily endorse the content provided by these organizations nor do the listed organizations necessarily endorse TG Therapeutics or its products.

### IMPORTANT SAFETY INFORMATION (CONT'D)

If you have any of the mentioned serious side effects during treatment with UKONIQ, your doctor may completely stop your treatment, stop your treatment for a period of time, or change your dose of UKONIQ.

**Before starting UKONIQ,** tell your healthcare provider about all of your medical conditions, including if you:

- Have an infection
- Have abdominal or intestinal problems
- Have liver problems
- Are allergic to FD&C Yellow No. 5 (tartrazine) or aspirin. UKONIQ tablets contain FD&C Yellow No. 5 (tartrazine), which may cause allergic-type reactions (including bronchial asthma) in certain people, especially people who also have an allergy to aspirin

## Important Information about UKONIQ

### What is UKONIQ?

**UKONIQ** is a prescription medicine used to treat adults with:

- **Marginal zone lymphoma (MZL)** when the disease has come back or did not respond to treatment and who have received at least 1 certain type of prior treatment
- **Follicular lymphoma (FL)** when the disease has come back or did not respond to treatment and who have received at least 3 prior treatments

It is not known if UKONIQ is safe and effective in children.

**UKONIQ** was approved through a faster FDA review process based on the percentage of patients whose tumor size shrank or disappeared after treatment. There is an ongoing study to confirm the benefit of UKONIQ for these uses.



### IMPORTANT SAFETY INFORMATION (CONT'D)

- Are pregnant or plan to become pregnant. UKONIQ can harm your unborn baby. If you are able to become pregnant, your healthcare provider will do a pregnancy test before starting treatment with UKONIQ.
  - Females and males with female partners who are able to become pregnant should use effective birth control (contraception) during treatment with UKONIQ and for 1 month after the last dose of UKONIQ. Talk to your healthcare provider about birth control methods that may be right for you. Tell your healthcare provider right away if you or your partner become pregnant or think you are pregnant during treatment with UKONIQ.
- Are breastfeeding or plan to breastfeed. It is not known if UKONIQ passes into your breast milk. Do not breastfeed during treatment with UKONIQ and for 1 month after the last dose of UKONIQ.

Please see Important Safety Information throughout and Important Facts about UKONIQ on pages 10-11.

**UKONIQ**<sup>™</sup>  
umbralisib 200 mg  
tablets

# Important Facts About UKONIQ

Read this brief summary carefully before taking UKONIQ. This summary does not take the place of talking to your healthcare provider about your medical condition or treatment.

## What is UKONIQ?

UKONIQ is a prescription medicine used to treat adults with:

- **Marginal zone lymphoma (MZL)** when the disease has come back or did not respond to treatment and who have received at least one certain type of prior treatment.
- **Follicular lymphoma (FL)** when the disease has come back or did not respond to treatment and who have received at least three prior treatments.

It is not known if UKONIQ is safe and effective in children.

UKONIQ was approved through a faster FDA review process based on the percentage of patients whose tumor size shrank or disappeared after treatment. There is an ongoing study to confirm the benefit of UKONIQ for these uses.

## What is the most important information I should know about UKONIQ?

**UKONIQ can cause serious side effects, including:**

**Infections.** UKONIQ can cause serious infections that may lead to death. Tell your healthcare provider right away if you have fever, chills, weakness, flu-like symptoms, or any other signs of infection during treatment with UKONIQ.

**Low white blood cell count (neutropenia).** Neutropenia is common with UKONIQ treatment and can sometimes be serious. Your healthcare provider will check your blood counts regularly during treatment with UKONIQ. Tell your healthcare provider right away if you have a fever or any signs of infection during treatment with UKONIQ.

**Diarrhea or inflammation of your intestine (colitis).** Diarrhea is common during UKONIQ treatment and can sometimes be serious. Tell your healthcare provider right away if you have diarrhea that does not go away or worsening diarrhea, stool with mucus or blood, or if you have severe stomach area (abdominal) pain during treatment with UKONIQ. Drink plenty of fluids during treatment with UKONIQ to help prevent dehydration from diarrhea.

**Liver problems.** Abnormal liver function blood test results are common and can sometimes be serious. Your healthcare provider will do blood tests before and during your treatment with UKONIQ to check for liver problems. Tell your healthcare provider right away if you have any of the following symptoms of liver problems: yellowing of your skin or the white part of your eyes (jaundice); dark or brown (tea colored) urine; pain in the upper right side of your stomach area (abdomen); bleeding or bruising more easily than normal

**Severe skin reactions.** Rashes and other skin reactions are common with UKONIQ treatment and can sometimes be severe and may lead to death. Tell your healthcare provider right away if you get a new or worsening skin rash or other signs of a severe skin reaction during treatment with UKONIQ, including: painful sores or ulcers on your skin, lips, or in your mouth; rash with blisters or peeling skin; rash with itching; rash with fever.

If you have any of the above serious side effects during treatment with UKONIQ, your doctor may completely stop your treatment, stop your treatment for a period of time, or change your dose of UKONIQ.

## Before starting UKONIQ, tell your healthcare provider about all of your medical conditions, including if you:

- have an infection
- have abdominal or intestinal problems
- have liver problems
- are allergic to FD&C Yellow No. 5 (tartrazine) or aspirin. UKONIQ tablets contain FD&C Yellow No. 5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain people, especially people who also have an allergy to aspirin
- are pregnant or plan to become pregnant. UKONIQ can harm your unborn baby. If you are able to become pregnant, your healthcare provider will do a pregnancy test before starting treatment with UKONIQ
  - Females who are able to become pregnant should use effective birth control (contraception) during treatment with UKONIQ and for 1 month after the last dose of UKONIQ. Talk to your healthcare provider about birth control methods that may be right for you. Tell your healthcare provider right away if you become pregnant or think you are pregnant during treatment with UKONIQ
  - Males with female partners who are able to become pregnant should use effective birth control (contraception) during treatment with UKONIQ and for 1 month after the last dose of UKONIQ
- are breastfeeding or plan to breastfeed. It is not known if UKONIQ passes into your breast milk. Do not breastfeed during treatment with UKONIQ and for 1 month after the last dose of UKONIQ

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

## What are the possible side effects of UKONIQ?

UKONIQ can cause serious side effects, including those in the “What is the most important information I should know about UKONIQ?” section.

The most common side effects of UKONIQ include changes in certain kidney function blood tests, tiredness, nausea, muscle or bone pain, low red blood cell count (anemia), low platelet count, upper respiratory tract infection, vomiting, abdominal pain, and decreased appetite.

These are not all the possible side effects with UKONIQ. Call your healthcare provider for medical advice about side effects.

This information provided here is not comprehensive. To get more information about UKONIQ, talk to your healthcare provider or pharmacist. You can also call TG Therapeutics at 1-877-848-9462 or visit UKONIQ.com to obtain the FDA-approved product labeling.

You are encouraged to report negative side effects or quality complaints of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

**UKONIQ**<sup>™</sup>  
umbralisib 200 mg  
tablets



For more information about UKONIQ, please visit [UKONIQ.com](http://UKONIQ.com)

## IMPORTANT SAFETY INFORMATION (CONT'D)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

### The most common side effects of UKONIQ include:

- Changes in certain kidney function blood tests
- Tiredness
- Nausea
- Muscle or bone pain
- Low red blood cell count (anemia)
- Low platelet count
- Upper respiratory tract infection
- Vomiting
- Abdominal pain
- Decreased appetite

These are not all the possible side effects with UNONIQ. Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see Important Safety Information throughout and Important Facts about UKONIQ on pages 10-11.



© 2021 TG Therapeutics, Inc. All rights reserved. UKONIQ is a trademark of TG Therapeutics, Inc. All other trademarks referenced herein are the property of their respective owners.  
05/2021 US-UMB-2000078

**UKONIQ**<sup>™</sup>  
umbralisib 200 mg  
tablets